

Monopoles and Knots in Skyrme Theory

Y. M. Cho

Department of Physics, College of Natural Sciences, Seoul National University, Seoul 151-742, Korea
ymcho@yongmin.snu.ac.kr

We show that the Skyrme theory actually is a theory of monopoles which allows a new type of solitons, the topological knots made of monopole-anti-monopole pair, which is different from the well-known skyrmions. Furthermore we derive a generalized Skyrme action from the Yang-Mills action of QCD, which we propose to be an effective action of QCD in the infra-red limit. We discuss the physical implications of our results.

PACS numbers: 21.60.Fw, 12.38.-t, 03.50.-z, 02.40.-k, 11.10.Lm

The proposal that the Skyrme theory could describe an effective theory of QCD in the low energy limit has become very popular [1,2]. In this view the skyrmions are interpreted as the baryons, and numerous evidences have been put forward which support this view [3]. Motivated by the success of the proposal many people have tried to derive the Skyrme action from QCD. But a concrete theoretical proof of the proposal, that one can actually derive the Skyrme action from QCD, has remained very difficult [4].

The purpose of this Letter is two-fold. First we show that the Skyrme theory actually has a much richer soliton spectrum. Indeed we show that it is a theory of self-interacting non-Abelian monopoles, and prove that it allows a new type of knotted solitons very similar to the topological knots in the Skyrme-Faddeev theory of non-linear sigma model [5]. The other purpose is to present a new evidence that the Skyrme theory is indeed very closely related to QCD. In fact, by reparametrizing the gauge potential, we derive a generalized Skyrme action from the Yang-Mills action of QCD which we propose to be an effective theory of QCD in the low energy limit. The assertion that the Skyrme theory could be derived from QCD is perhaps not suprising. But the fact that the Skyrme theory has a new type of solitons is really unexpected, which put the Skyrme theory in a completely new perspective.

Let us start from the Skyrme theory. With

$$U = \exp(\xi \frac{\vec{\sigma}}{2i} \cdot \hat{n}) = \cos \frac{\xi}{2} - i(\vec{\sigma} \cdot \hat{n}) \sin \frac{\xi}{2},$$

$$L_\mu = U \partial_\mu U^\dagger, \quad (\hat{n}^2 = 1) \quad (1)$$

the Skyrme Lagrangian is expressed as

$$\begin{aligned} \mathcal{L} &= \frac{\mu^2}{4} \text{tr} L_\mu^2 + \frac{\alpha}{32} \text{tr} ([L_\mu, L_\nu])^2 \\ &= -\frac{\mu^2}{4} \left[\frac{1}{2} (\partial_\mu \xi)^2 + (1 - \cos \xi) (\partial_\mu \hat{n})^2 \right] \\ &\quad - \frac{\alpha}{16} \left[\frac{1 - \cos \xi}{2} (\partial_\mu \xi \partial_\nu \hat{n} - \partial_\nu \xi \partial_\mu \hat{n})^2 \right. \end{aligned}$$

$$\left. + (1 - \cos \xi)^2 (\partial_\mu \hat{n} \times \partial_\nu \hat{n})^2 \right]. \quad (2)$$

The equation of motion is given by

$$\begin{aligned} &\frac{\mu^2}{4} [\partial^2 \xi - \sin \xi (\partial_\mu \hat{n})^2] \\ &+ \frac{\alpha}{32} \sin \xi (\partial_\mu \xi \partial_\nu \hat{n} - \partial_\nu \xi \partial_\mu \hat{n})^2 \\ &+ \frac{\alpha}{8} (1 - \cos \xi) \partial_\mu [(\partial_\mu \xi \partial_\nu \hat{n} - \partial_\nu \xi \partial_\mu \hat{n}) \cdot \partial_\nu \hat{n}] \\ &- \frac{\alpha}{8} (1 - \cos \xi) \sin \xi (\partial_\mu \hat{n} \times \partial_\nu \hat{n})^2 = 0, \\ &\partial_\mu \left\{ \frac{\mu^2}{4} (1 - \cos \xi) \hat{n} \times \partial_\mu \hat{n} \right. \\ &+ \frac{\alpha}{16} (1 - \cos \xi) [(\partial_\nu \xi)^2 \hat{n} \times \partial_\mu \hat{n} - (\partial_\mu \xi \partial_\nu \xi) \hat{n} \times \partial_\nu \hat{n}] \\ &\left. + \frac{\alpha}{8} (1 - \cos \xi)^2 (\hat{n} \cdot \partial_\mu \hat{n} \times \partial_\nu \hat{n}) \partial_\nu \hat{n} \right\} = 0. \end{aligned} \quad (3)$$

With the spherically symmetric ansatz

$$\xi = \xi(r), \quad \hat{n} = \hat{r}, \quad (4)$$

(3) is reduced to

$$\begin{aligned} &\frac{d^2 \xi}{dr^2} + \frac{2}{r} \frac{d\xi}{dr} - \frac{2 \sin \xi}{r^2} - \frac{\alpha}{\mu^2} \left[\frac{\sin^2(\xi/2)}{r^2} \frac{d^2 \xi}{dr^2} \right. \\ &\left. + \frac{\sin \xi}{2r^2} \left(\frac{d\xi}{dr} \right)^2 - \frac{2 \sin \xi \sin^2(\xi/2)}{r^4} \right] = 0. \end{aligned} \quad (5)$$

Imposing the boundary condition $\xi(0) = 2\pi$ and $\xi(\infty) = 0$, one has the well-known skyrmions. The reason for the solitons, of course, is the non-trivial homopopy $\pi_3(S^3)$ defined by (1) [1,2].

Now, we show that (3) actually allows a new type of knot solitons. To see this notice that with

$$\xi = \pi, \quad (6)$$

(3) is reduced to the following equation for $\hat{n}$

$$\hat{n} \times \partial^2 \hat{n} + \frac{\alpha}{\mu^2} (\partial_\mu H_{\mu\nu}) \partial_\nu \hat{n} = 0, \quad (7)$$

where

$$H_{\mu\nu} = \hat{n} \cdot (\partial_\mu \hat{n} \times \partial_\nu \hat{n}) = \partial_\mu C_\nu - \partial_\nu C_\mu.$$

Notice that since $H_{\mu\nu}$ is closed, one can always introduce the potential C_μ for the field $H_{\mu\nu}$, as far as $\hat{n}$ is smooth everywhere. But (7) is precisely the knot equation that we obtain from the Skyrme- Faddeev action [6]

$$\mathcal{L}_{SF} = -\frac{\mu^2}{2}(\partial_\mu \hat{n})^2 - \frac{\alpha}{4}(\partial_\mu \hat{n} \times \partial_\nu \hat{n})^2. \quad (8)$$

This immediately tells that the knots in this theory automatically become the solutions of the Skyrme theory. This guarantees the existence of the knots in Skyrme theory. This is striking. But considering the fact that the Skyrme action (1) is indeed an obvious generalization of the Skyrme-Faddeev action (8), one should not be too suprised at the knots.

Another remarkable point is that the Skyrme theory actually has the non-Abelian monopoles as the classical solution. This must be clear because the knot equation (7) allows the monopoles [6,7]. Indeed it is clear that

$$\xi = \pi, \quad \hat{n} = \hat{r}, \quad (9)$$

forms a solution of (3) except at the origin, because

$$\partial^2 \hat{r} = -\frac{2}{r^2} \hat{r}, \quad \partial_\mu H_{\mu\nu} = 0. \quad (10)$$

Notice that (9) forms a solution of (3) even without the non-linear interaction (i.e., with $\alpha = 0$). As we will see soon the solution can naturally be identified as the monopole [6,7].

It is really remarkable that the Skyrme theory has such a rich spectrum of classical solutions, all of which are of topological origin. Skyrme noticed that the existence of $\pi_3(S^3)$ described by (1) produces the topological skyrmions. But our analysis tells that the topological field $\hat{n}$ defines two more homotopies $\pi_3(S^2)$ and $\pi_2(S^2)$, which produce their own knots and monopoles.

At this point we may ask what is the relationship among these topological solutions. To see this remember that the baryonic charge of the skyrmions is given by

$$Q_s = \frac{1}{24\pi^2} \int \epsilon_{ijk} \text{tr} (L_i L_j L_k) \\ = \frac{1}{16\pi^2} \int \epsilon_{ijk} (1 - \cos \xi) H_{ij} \partial_k \xi d^3 x. \quad (11)$$

On the other hand the knot and monopole charges are given by [6,7]

$$Q_k = \frac{1}{32\pi^2} \int \epsilon_{ijk} C_i H_{jk} d^3 x, \\ Q_m = \frac{1}{8\pi} \int \epsilon_{ijk} H_{ij} d\sigma_k, \quad (12)$$

where C_μ is the potential of $H_{\mu\nu}$ defined by (7). From this we deduce the followings. First the skyrmions carry the monopole charge of the topological field $\hat{n}$. This suggests that the skyrmions are actually monopoles whose energy is made finite by the non-linear interaction with the scalar field ξ . More importantly, with the Hopf fibering of $S^3 \simeq S^2 \times S^1$ of the $SU(2)$ space, ξ adds the $U(1)$ charge of the fibre S^1 to the monopoles, which makes the baryonic charge different from the monopole charge. Secondly the knots carry neither the baryonic nor monopole charges (At the same time the skyrmions and the monopoles have no knot charge, because the potential of their magnetic field become singular). This suggests that the knots should be interpreted as the “mesons”, or more precisely the “glueballs”, made of the monopole-anti-monopole pairs. The non-linear self-interaction of $\hat{n}$ not only makes their energy finite, but also provides the topological stability to the knots.

This picture helps us to clarify the physical content of Skyrme theory. Skyrme suggested that it is probably a theory of non-linear sigma field, which allows the topological skyrmions. But our analysis suggests a different interpretation. *The Skyrme theory should really be viewed as a theory of the monopole field $\hat{n}$ interacting with the scalar field ξ . The non-linear self-interaction of $\hat{n}$ produces the knots made of monopole-anti-monopole pairs, and ξ adds the $U(1)$ charge of the Hopf fibre to the monopoles to produce the topological skyrmions.*

Now, we establish a deep connection between Skyrme action and QCD. For this we need a better understanding of QCD. A unique feature of the non-Abelian gauge theory is its non-trivial topology, and it is essential to take care of these topological characters in the dynamics for us to understand QCD. For this purpose it is very important to reparametrize the non-Abelian gauge potential in terms of the restricted potential of the maximal Abelian subgroup H of the gauge group G and the gauge covariant valence potential of the remaining G/H degrees [8,9]. To see this consider $SU(2)$ for simplicity. A natural way to take care of the topological degrees into the dynamics is to introduce a topological field $\hat{n}$ which selects the color direction at each space-time point, and to decompose the non-Abelian connection into the restricted potential which makes $\hat{n}$ a covariant constant and the valence potential which forms a covariant vector field [8,9],

$$\vec{A}_\mu = A_\mu \hat{n} - \frac{1}{g} \hat{n} \times \partial_\mu \hat{n} + \vec{X}_\mu = \hat{A}_\mu + \vec{X}_\mu, \\ (A_\mu = \hat{n} \cdot \vec{A}_\mu, \quad \hat{n}^2 = 1, \quad \hat{n} \cdot \vec{X}_\mu = 0). \quad (13)$$

Notice that the restricted potential $\hat{A}_\mu$ is precisely the connection which leaves $\hat{n}$ invariant under the parallel transport,

$$\hat{D}_\mu \hat{n} = \partial_\mu \hat{n} + g \hat{A}_\mu \times \hat{n} = 0. \quad (14)$$

Under the infinitesimal gauge transformation

$$\delta \hat{n} = -\vec{\alpha} \times \hat{n}, \quad \delta \vec{A}_\mu = \frac{1}{g} D_\mu \vec{\alpha}, \quad (15)$$

one has [8,9]

$$\begin{aligned} \delta A_\mu &= \frac{1}{g} \hat{n} \cdot \partial_\mu \vec{\alpha}, \quad \delta \hat{A}_\mu = \frac{1}{g} \hat{D}_\mu \vec{\alpha}, \\ \delta \vec{X}_\mu &= -\vec{\alpha} \times \vec{X}_\mu. \end{aligned} \quad (16)$$

This shows that $\hat{A}_\mu$ by itself describes an $SU(2)$ connection which enjoys the full $SU(2)$ gauge degrees of freedom. Furthermore the valence potential $\vec{X}_\mu$ transforms covariantly under the gauge transformation. More importantly the decomposition is gauge-independent. Once the gauge covariant topological field $\hat{n}$ is given, the decomposition follows automatically independent of the choice of a gauge.

This decomposition, which has recently become known as the ‘‘Cho decomposition’’ [10] or the ‘‘Cho-Faddeev-Niemi decomposition’’ [11], was introduced long time ago in an attempt to demonstrate the monopole condensation in QCD [8,9]. But only recently the importance of the decomposition in clarifying the non-Abelian dynamics has become appreciated by many authors [10,11]. Indeed it is this decomposition which has played a crucial role to establish the Abelian dominance in the Wilson loops in QCD [12], and the possible connection between the Skyrme-Faddeev action and the effective action of QCD [13].

An important feature of $\hat{A}_\mu$ is that, as an $SU(2)$ potential, it retains the full topological characteristics of the original non-Abelian potential. This is due to the topological field $\hat{n}$ in it. Clearly the isolated singularities of $\hat{n}$ defines $\pi_2(S^2)$ which describes the non-Abelian monopoles. Indeed $\hat{A}_\mu$ with $A_\mu = 0$ and $\hat{n} = \hat{r}$ describes precisely the Wu-Yang monopole [7,8]. So one can identify the non-Abelian magnetic potential by

$$\vec{C}_\mu = -\frac{1}{g} \hat{n} \times \partial_\mu \hat{n}, \quad (17)$$

which generates the magnetic field

$$\begin{aligned} \vec{H}_{\mu\nu} &= \partial_\mu \vec{C}_\nu - \partial_\nu \vec{C}_\mu + g \vec{C}_\mu \times \vec{C}_\nu = -\frac{1}{g} \partial_\mu \hat{n} \times \partial_\nu \hat{n} \\ &= -\frac{1}{g} (\partial_\mu C_\nu - \partial_\nu C_\mu) \hat{n}. \end{aligned} \quad (18)$$

Furthermore, with the S^3 compactification of R^3 , $\hat{n}$ characterizes the Hopf invariant $\pi_3(S^2)$ which describes the topologically distinct vacua [6,14]. So the restricted potential inherits all the essential topological properties of the non-Abelian gauge theory.

The decomposition (13) leads us to establish the common ground between QCD and Skyrme theory, and to clarify the meaning of the Skyrme theory. To see this notice that with (17) and (18) the Skyrme-Faddeev Lagrangian (8) can be expressed as

$$\mathcal{L}_{SF} = -\frac{\alpha}{4} \vec{H}_{\mu\nu}^2 - \frac{\mu^2}{2} \vec{C}_\mu^2. \quad (19)$$

This allows us to interpret the solution (9) as the monopoles, and to translate the vacuum number of QCD to the knot quantum number of Skyrme theory [6,14].

With the decomposition (13) we have

$$\vec{F}_{\mu\nu} = \hat{F}_{\mu\nu} + \hat{D}_\mu \vec{X}_\nu - \hat{D}_\nu \vec{X}_\mu + g \vec{X}_\mu \times \vec{X}_\nu, \quad (20)$$

so that the Yang-Mills Lagrangian could be expressed as

$$\begin{aligned} \mathcal{L} &= -\frac{1}{4} \hat{F}_{\mu\nu}^2 - \frac{\mu^2}{2} \hat{A}_\mu^2 - \frac{1}{4} (\hat{D}_\mu \vec{X}_\nu - \hat{D}_\nu \vec{X}_\mu)^2 \\ &\quad - \frac{g}{2} \hat{F}_{\mu\nu} \cdot (\vec{X}_\mu \times \vec{X}_\nu) - \frac{g^2}{4} (\vec{X}_\mu \times \vec{X}_\nu)^2, \end{aligned} \quad (21)$$

where we have inserted a mass term for the restricted potential $\hat{A}_\mu$. One may object the introduction of the mass term, because it will certainly violate the gauge invariance of QCD. But notice that, after the dynamical symmetry breaking with the monopole condensation, one could assume (simple mindedly) that the restricted potential acquires a mass [13,15]. This would justify the introduction of the mass term in the infra-red limit of QCD.

With this preparation we now derive a generalized Skyrme theory from the Lagrangian (21). To do this let

$$\vec{X}_\mu = f_1 \partial_\mu \hat{n} + f_2 \hat{n} \times \partial_\mu \hat{n}, \quad (22)$$

and find that the Lagrangian is expressed as

$$\begin{aligned} \mathcal{L} &= -\frac{1}{4} [F_{\mu\nu} + (\phi^* \phi - \frac{1}{g})(\hat{n} \cdot \partial_\mu \hat{n} \times \partial_\nu \hat{n})]^2 \\ &\quad - \frac{i}{2} (D_\mu \phi)^* (D_\nu \phi) (\hat{n} \cdot \partial_\mu \hat{n} \times \partial_\nu \hat{n}) \\ &\quad - \frac{1}{4} |D_\mu \phi \partial_\nu \hat{n} - D_\nu \phi \partial_\mu \hat{n}|^2 - \frac{\mu^2}{2} [A_\mu^2 + \frac{1}{g^2} (\partial_\mu \hat{n})^2], \end{aligned} \quad (23)$$

where

$$\phi = f_1 + i f_2, \quad D_\mu \phi = (\partial_\mu + i g A_\mu) \phi.$$

Furthermore, with $\partial_\mu \phi = 0$ and $A_\mu = \partial_\mu \xi$, we have

$$\begin{aligned} \mathcal{L} &= -\frac{1}{4g^2} (1 - g \phi^* \phi)^2 (\partial_\mu \hat{n} \times \partial_\nu \hat{n})^2 \\ &\quad - \frac{1}{4} \phi^* \phi (\partial_\mu \xi \partial_\nu \hat{n} - \partial_\nu \xi \partial_\mu \hat{n})^2 \\ &\quad - \frac{\mu^2}{2} [(\partial_\mu \xi)^2 + \frac{1}{g^2} (\partial_\mu \hat{n})^2]. \end{aligned} \quad (24)$$

At this point a striking similarity between the Skyrme Lagrangian (1) and the QCD Lagrangian (23) becomes unmistakable. Of course (23) is different from (1). But notice that it has a definite advantage over (1), in the sense that it comes directly from QCD. With this observation we can propose (23) as a generalized Skyrme Lagrangian which could replace (1) as an effective theory of QCD in the low energy limit.

We close with the following comments.

1) The assertion that one can derive the Skyrme action from QCD is understandable. Recently we have been able to derive a generalized Skyrme-Faddeev action from the one-loop effective action of QCD which we obtained with the monopole background [13,15]. This derivation was different from our derivation of the generalized Skyrme action discussed here. Our discussion here is based on a reparametrization of the potential and does not appear to resort to an effective action of QCD (except the insertion of the mass term of the restricted potential). But clearly the two results are consistent with each other. Indeed, as far as the generalized Skyrme theory could be viewed as a generalized Skyrme-Faddeev theory, one could say that the derivation of the generalized Skyrme-Faddeev action from the effective action of QCD [13,15] provides another evidence which supports our derivation in this paper.

2) Our result suggests that the Skyrme theory is a theory of monopoles interacting with the scalar field ξ . But here we like to argue that it is also a theory of confinement which confines the monopoles. To see this notice that in spite of the existence of the mass scale μ in the theory, the theory admits a long range monopole solutions. Furthermore, the mass scale does not describe the mass of the elementary fields of the theory. It describes the energy scale of the solitons. The classical monopole solutions carry an infinite energy and thus an infinite mass, while the monopole-anti-monopole bound states carry finite energy. This suggests that the theory, just like the Skyrme-Faddeev theory, is a theory of confinement [6]. What is remarkable is that the confinement is manifest already at the classical level. In QCD one has to produce a mass scale to prove the confinement. But here we already have the mass scale at the classical level which we do not have to produce from quantum correction.

3) We have proposed the generalized Skyrme action (23), instead of the Skyrme action (1), as an effective action of QCD in the low energy limit. In spite of the obvious differences they have a striking similarity. One common aspect is that both admit the knot solitons [16], which we have interpreted as the “glueballs”. We emphasize, however, that they are the magnetic glueballs which carry the color magnetic flux, while the glueballs in QCD (if they exist) are supposed to carry the color electric flux. This tells that our solitons should describe very exotic states. But the existence of such magnetic glueballs has already been predicted long time ago [8,9]. Once the monopole

condensation sets in, one should expect the fluctuation of the condensed vacuum. But obviously the fluctuation modes have to be magnetic, which could be identified as the magnetic glueballs (A new feature here is that they have a topological stability. This could be an artifact of the effective theory, not a genuine feature of QCD). If so, the remaining task is to look for a convincing experimental evidence of the magnetic glueball states in hadron spectrum [8,9].

The mass of the lightest knot in Skyrme theory can be estimated to be about 1.42 times the mass of the nucleon. A detailed discussion will be presented elsewhere [16].

ACKNOWLEDGEMENT

We thank Professor C. N. Yang for the illuminating discussions. The work is supported in part by a Korea Research Foundation Grant (KRF-2000-015-BP0072), and by the BK21 project of the Ministry of Education.

-
- [1] T. H. R. Skyrme, Proc. Roy. Soc. (London) **260**, 127 (1961); **262**, 237 (1961); Nucl. Phys. **31**, 556 (1962).
 - [2] See, for example, I. Zahed and G. Brown, Phys. Rep. **142**, 1 (1986), and the references therein.
 - [3] A. Jackson and M. Rho, Phys. Rev. Lett. **51**, 751 (1983); G. Atkins, C. Nappi, and E. Witten, Nucl. Phys. **B228**, 552 (1983).
 - [4] P. Simic, Phys. Rev. Lett. **55**, 40 (1985).
 - [5] L. Faddeev and A. Niemi, Nature **387**, 58 (1997); R. Battye and P. Sutcliffe, Phys. Rev. Lett. **81**, 4798 (1998).
 - [6] W. S. Bae, Y. M. Cho, and S. W. Kimm, hep-th/0105163, Phys. Rev. **D**, in press.
 - [7] Y. M. Cho, Phys. Rev. Lett. **44**, 1115 (1980); Phys. Lett. **B115**, 125 (1982); Y. M. Cho and D. Maison, Phys. Lett. **B391**, 360 (1997).
 - [8] Y. M. Cho, Phys. Rev. **D21**, 1080 (1980); J. Korean Phys. Soc. **17**, 266 (1984).
 - [9] Y. M. Cho, Phys. Rev. Lett. **46**, 302 (1981); Phys. Rev. **D23**, 2415 (1981).
 - [10] L. Faddeev and A. Niemi, Phys. Rev. Lett. **82**, 1624 (1999); Phys. Lett. **B449**, 214 (1999); **B464**, 90 (1999).
 - [11] E. Langman and A. Niemi, Phys. Lett. **B463**, 252 (1999); S. Shabanov, Phys. Lett. **B458**, 322 (1999); **B463**, 263 (1999); H. Gies, hep-th/0102026.
 - [12] Y. M. Cho, Phys. Rev. **D62**, 074009-1 (2000).
 - [13] Y. M. Cho and D. G. Pak, J. Korean Phys. Soc. **38**, 151 (2001); Y. M. Cho, H. W. Lee, and D. G. Pak, hep-th/9905215, Phys. Lett. **B**, in press.
 - [14] Y. M. Cho, Phys. Lett. **B81**, 25 (1979); P. van Vaal and A. Wipf, Phys. Lett. **B515**, 181 (2001).
 - [15] Y. M. Cho and D. G. Pak, hep-th/0006051, submitted to Phys. Rev. **D**.
 - [16] Y. M. Cho, and S. W. Kimm, to be published.